

DATA NOTE

Open Access



# 2020-2021 field seasons of Maize GxE project within the Genomes to Fields Initiative

Dayane Cristina Lima<sup>1\*</sup>, Alejandro Castro Aviles<sup>1</sup>, Ryan Timothy Alpers<sup>1</sup>, Alden Perkins<sup>1</sup>, Dylan L Schoemaker<sup>1</sup>, Martin Costa<sup>1</sup>, Kathryn J. Michel<sup>1</sup>, Shawn Kaeppler<sup>1</sup>, David Ertl<sup>2</sup>, Maria Cinta Romay<sup>3</sup>, Joseph L. Gage<sup>4</sup>, James Holland<sup>5</sup>, Timothy Beissinger<sup>6</sup>, Martin Bohn<sup>7</sup>, Edward Buckler<sup>8</sup>, Jode Edwards<sup>9</sup>, Sherry Flint-Garcia<sup>10</sup>, Michael A. Gore<sup>11</sup>, Candice N. Hirsch<sup>12</sup>, Joseph E. Knoll<sup>13</sup>, John McKay<sup>14</sup>, Richard Minyo<sup>15</sup>, Seth C. Murray<sup>16</sup>, James Schnable<sup>17</sup>, Rajandeep S. Sekhon<sup>18</sup>, Maninder P. Singh<sup>19</sup>, Erin E. Sparks<sup>20</sup>, Peter Thomison<sup>21</sup>, Addie Thompson<sup>19</sup>, Mitchell Tuinstra<sup>22</sup>, Jason Wallace<sup>23</sup>, Jacob D. Washburn<sup>10</sup>, Teclamariam Weldekidan<sup>20</sup>, Wenwei Xu<sup>24</sup> and Natalia de Leon<sup>1</sup>

## Abstract

**Objectives** This release note describes the Maize GxE project datasets within the Genomes to Fields (G2F) Initiative. The Maize GxE project aims to understand genotype by environment (GxE) interactions and use the information collected to improve resource allocation efficiency and increase genotype predictability and stability, particularly in scenarios of variable environmental patterns. Hybrids and inbreds are evaluated across multiple environments and phenotypic, genotypic, environmental, and metadata information are made publicly available.

**Data description** The datasets include phenotypic data of the hybrids and inbreds evaluated in 30 locations across the US and one location in Germany in 2020 and 2021, soil and climatic measurements and metadata information for all environments (combination of year and location), ReadMe, and description files for each data type. A set of common hybrids is present in each environment to connect with previous evaluations. Each environment had a collaborator responsible for collecting and submitting the data, the GxE coordination team combined all the collected information and removed obvious erroneous data. Collaborators received the combined data to use, verify and declare that the data generated in their own environments was accurate. Combined data is released to the public with minimal filtering to maintain fidelity to the original data.

**Keywords** Maize, Genotype by Environment, Grain Yield, Stability, Prediction

\*Correspondence:  
Dayane Cristina Lima  
dclima@wisc.edu

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Objective

The release of this data provides a unique resource to understand and dissect genotype-by-environment interactions in maize (*Zea mays* subsp. *mays* L.). Collaborators generate phenotypic, environmental, and metadata datasets to support a more comprehensive understanding of the opportunities and challenges associated with maize production in various environments. The Maize GxE project data is made available to the public in its original form, with minimum filtering to remove erroneous data or as specified by collaborators and in the description files. This approach ensures that the publicly available data contains the maximum amount of information collected by project collaborators and empowers users to define their quality controls based on their specific goals.

A set of 1184 publicly available hybrids were evaluated in the 2020 and 2021 seasons across 30 different locations. The main group of hybrids was produced by the cross of doubled-haploid (DH) inbred lines from the Wisconsin Stiff Stalk MAGIC population (WI-SS-MAGIC), crossed with three ex-PVP inbred testers, PHZ51, PHP02, and PHK76 [1]. The WI-SS-MAGIC population involves the inbreds B73, B84, NKH8431, LH145, PHB47, and PHJ40 as parents in the initial crosses, and a detailed description of the population creation and DH production is in Michel et al. (2022) [1]. The testers were selected to allow adaptation of materials to the wide array of maturities sampled across the project. Inbred tester PHZ51 was used in southern locations (DEH1, GAH1, GAH2, IAH1, IAH2, IAH3, IAH4, MOH1, NCH1, NEH1, NEH2, NEH3, NYH3, SCH1, TXH1, TXH2, TXH3, WIH2), PHK76 in the Midwest and intermediate locations (DEH1, IAH1, IAH2, IAH3, IAH4, ILH1, INH1, MOH1, NCH1, NYH3, WIH2), and PHP02 in the northern locations (DEH1, GEH1, IAH2, IAH3, IAH4, MIH1, MNH1, MOH1, NYH2, OHH1, WIH1, WIH2, WIH3). Six locations (mega-locations) had hybrids created using all three testers (DEH1, IAH2, IAH3, IAH4, MOH1, WIH2). Additional smaller-scale experiments were conducted alongside the main experiment for additional phenotyping and/or deployment of novel phenotyping methods and tools across approximately 83% of the locations. These experiments included the external Yellow Stripes (YS), same set of check hybrids known as 'Yellow Stripe' used to connect location and years, but evaluated in a different experiment; the High-Intensity Phenotyping Site (HIPS), which tested 22 hybrids (HIP\_Hybrid) and 22 inbreds (HIP\_Inbred). HIPS was introduced in 2020 as a more comprehensive phenotyping set for aerial high-throughput phenotyping platforms. The choice of hybrids and inbreds was based on their historical importance and relevance to other connected projects.

## Data description

The 2020 and 2021 datasets are publicly available via CyVerse/iPlant. These datasets contain phenotypic, environmental, soil, and supplemental data, and have been structured according to the specifications outlined in Table 1.

- **Phenotypic data:** Phenotypic measurements that follow a standard set of instructions, available at [genomes2fields.org](https://genomes2fields.org). Standard traits include days to anthesis, days to silking, ear height, plant height, stand count, stalk lodging, root lodging, grain moisture, test weight, plot weight, and grain yield. Both raw data and minimally quality-controlled (clean) data are reported separately. Out of range observations were set to missing values following the rules described in the readMe files.
- **Environmental dataset:** WatchDog 2700 weather stations (Spectrum Technologies) were placed at each field site. Data were collected at 30-min intervals, or according with collaborator set up, from planting through harvest at each location. The geographic locations of the experiments are not identical across years due to crop rotation management practices; thus, the locations of the weather stations vary across years. Each station measured wind speed, direction, and gust; air temperature, dewpoint, relative humidity; soil temperature and moisture; rainfall and solar radiation. Additional measurements taken at selected sites included soil electrical conductivity, ultraviolet light, carbon dioxide, and photosynthetically active radiation. Instructions for weather station maintenance activities including pre-season tasks, field setup, maintenance throughout the growing season, and removal are available on the G2F webpage [2].
- **Soil dataset:** Each field location collected soil samples that represent the experiment field according to the instructions available on the G2F webpage.
- **Supplemental dataset:** Supplemental information consists of metadata (any field-level data collected at planting, in season, and/or at harvest), agronomic information (list of products, nutrients, and irrigation applied), and cooperator list (collaborators responsible for the field locations in 2020 and 2021).

## Limitations

These datasets contain missing data. Missing data includes data not reported by collaborators or erroneous data as specified on the readMe and description files. Genotypic data is not included in this release. Locations that did not collect data due to the complete loss of the experiment are listed on the cooperator list.

**Table 1** Overview of 2020 and 2021 planting seasons datasets

Label	Name of data file/data set	File types (Extension)	Data repository and identifier
Data file 1	readme.txt	.txt	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 2	g2f_2020_phenotypic_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 3	g2f_2020_phenotypic_raw_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 4	g2f_2020_phenotypic_clean_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 5	g2f_2020_weather_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 6	g2f_2020_weather_readMe.txt	.txt	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 7	2020_weather_raw.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 8	2020_weather_cleaned.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 9	g2f_2020_soil_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 10	g2f_2020_soil_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 11	g2f_2020_supplemental_information.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 12	g2f_2020_agronomic_information.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 13	g2f_2020_cooperators_list.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 14	g2f_2020_field_metadata.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/hzss-a865">https://doi.org/10.25739/hzss-a865</a> ) [3]
Data file 15	readme.txt	.txt	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 16	g2f_2021_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 17	g2f_2021_phenotypic_raw_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 18	g2f_2021_phenotypic_clean_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 19	g2f_2021_weather_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 20	g2f_2021_weather_readMe.txt	.txt	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 21	g2f_2021_weather_raw.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 22	g2f_2021_weather_cleaned.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 23	g2f_2021_soil_data_description.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 24	g2f_2021_soil_data.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 25	g2f_2021_supplemental_information.pdf	.pdf	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 26	g2f_2021_agronomic_information.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 27	g2f_2021_cooperators_list.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]
Data file 28	g2f_2021_field_metadata.csv	.csv	CyVerse ( <a href="https://doi.org/10.25739/5ae3-sw62">https://doi.org/10.25739/5ae3-sw62</a> ) [4]

**Abbreviations**

G2F	Genomes to Fields
DH	doubled-haploid
GxE	Genotype by environment

**Acknowledgements**

We gratefully acknowledge contributions from many field managers, crop research coordinators, staff, graduate students, student interns and data collectors including: Dustin Eilert, Marina Borsecnik, Renata Barcelos and Ben Fischer (de Leon/Kaeppler labs, University of Wisconsin - Madison). Naomi Rodman, Hadden Duling, Nathan Moore, Evonne Pinto, Marielle Quinn, & Kai Sanders (Wallace lab). Jason Brewer (USDA-ARS, Raleigh, NC). Trevor Perla, Paige Coffee, Will Deems, and Amy Deariso (ARS-Tifton, GA). Brian Hearn, Gunnar Isaacs, Buddy Willey, Dheeraj Danthuluri, and Raphael Kim (University of Delaware). Tom Siler (Michigan State University). Katherine Guill, Brady Blanton, Mia Ruppel, Grace Sidberry, Daniel Kick (Washburn lab, USDA-ARS). Amanda Gilbert, Peter Hermanson, and Thomas Hoverstad (University of Minnesota).

**Authors' contributions**

Data management team: DCL, ACA, RTA, AP, DLS, MC, JH. Data contributors: DCL, ACA, RTA, NdL, SK, MCR, JLG, JH, TB, MB, EB, JE, SFG, MAG, CNH, JEK, JM, RM, SCM, JS, RSS, MPS, EES, PT, AT, MT, JW, JDW, TW, WX. Communication: NdL, DE, SK. The data management team aggregated and curated data resources and made them available to other researchers. Data contributors were responsible for advising on data collection methods, collecting the data, reviewing data collection and curation methods, and the resulting datasets. Communicating authors guided the data management process, including data collection, curation, and distribution, and oversaw the writing of the manuscript. All authors reviewed and contributed to the manuscript and agreed with publication.

**Funding**

We gratefully acknowledge support from: National Corn Growers Association, Iowa Corn Promotion Board, Georgia Corn Commission, Nebraska Corn Board, Ohio Corn Marketing Program, Corn Marketing Program of Michigan, Texas Corn Producers Board, University of Göttingen startup funds, USDA-ARS, and USDA Germplasm Enhancement of Maize program.

**Data availability**

The data described in this Data note can be freely and openly accessed on CyVerse at <https://doi.org/10.25739/hzss-a865> (2020 Field Season [3]) and <https://doi.org/10.25739/5ae3-sw62> (2021 Field Season [4]).

**Declarations****Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare no competing interests.

**Author details**

<sup>1</sup>Department of Agronomy, University of Wisconsin - Madison, Madison, WI 53706, USA

<sup>2</sup>Iowa Corn Promotion Board, Johnston, IA 50131, USA

<sup>3</sup>Institute for Genomic Diversity, Cornell University, Ithaca, NY 14853, USA

<sup>4</sup>Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC 27695, USA

<sup>5</sup>USDA-ARS Plant Science Research Unit, Raleigh, NC 27606, USA

<sup>6</sup>Department of Crop Science, Center for Integrated Breeding Research, University of Göttingen, Carl-Neuberg-Weg 1, 37075 Göttingen, Germany,

<sup>7</sup>University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

<sup>8</sup>USDA-ARS and Cornell University, Ithaca, NY 14853, USA

<sup>9</sup>USDA ARS CICGRU, 716 Farmhouse Ln, Ames, IA 50011-1051, USA

<sup>10</sup>USDA-ARS, Plant Genetics Research Unit, University of Missouri, 205 Curtis Hall, Columbia, MO 65211, USA

<sup>11</sup>Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University, Ithaca, NY 14853, USA

<sup>12</sup>Department of Agronomy and Plant Genetics, University of Minnesota, St Paul, MN 55108, USA

<sup>13</sup>USDA-ARS Crop Genetics and Breeding Research Unit, Tifton, GA 31793, USA

<sup>14</sup>Department of Agricultural Biology, Colorado State University, Fort Collins, CO 80523, USA

<sup>15</sup>Department of Horticulture and Crop Science, College of Food, Agricultural, and Environmental Sciences, Ohio State University, Wooster, OH 44691, USA

<sup>16</sup>Department of Soil and Crop Sciences, Texas A&M University, College Station, TX 77843, USA

<sup>17</sup>Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, NE 68588, USA

<sup>18</sup>Department of Genetics and Biochemistry, Clemson University, Clemson, SC 29634, USA

<sup>19</sup>Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, MI 48824, USA

<sup>20</sup>Department of Plant and Soil Sciences, University of Delaware, Newark, DE 19716, USA

<sup>21</sup>Ohio State University, Columbus, OH 43210, USA

<sup>22</sup>Department of Agronomy, Purdue University, West Lafayette, IN 47907, USA

<sup>23</sup>Department of Crop & Soil Sciences, University of Georgia, Athens, GA 30602, USA

<sup>24</sup>Texas A&M University, College Station, TX 77843, USA

Received: 8 May 2023 / Accepted: 17 July 2023

Published online: 14 September 2023

## References

1. Michel KJ, Lima DC, Hundley H, Singan V, Yoshinaga Y, Daum C et al. Genetic mapping and prediction of flowering time and plant height in a maize Stiff Stalk MAGIC population. *Genetics*. 2022;221.
2. Genomes to Fields resources. 2023. <https://www.genomes2fields.org/resources/>.
3. G2F Consortium. Genomes to Fields 2020 dataset. CyVerse Data Commons. 2020. <https://doi.org/10.25739/hzsz-a865>.
4. G2F Consortium. Genomes to Fields 2021 dataset. CyVerse Data Commons. 2021. <https://doi.org/10.25739/5ae3-sw62>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.